



Consider an object that moves along a horizontal frictionless surface (*e.g.*, an air hockey puck on a level air table). Suppose that the object moves under the influence of a net force expressed as follows:

$$\vec{F}_{net}(x, y) = (-k_x x \hat{i}) + (-k_y y \hat{j})$$

Note: The above net force can be modeled by two long springs connecting the air hockey puck to two edges of the air table. One spring, with force constant k_x , would be oriented in one direction; the other spring with force constant k_y , would be oriented perpendicular to the first spring.

Each diagram below corresponds to a specific experiment. The relative values of the force constants k_x and k_y and the initial conditions of the motion are given in each case.


For each case below, carefully sketch a qualitatively correct x - y trajectory that the object might follow. Explain the reasoning you used to decide how to draw the trajectory for each case.

<p>a. The force constants are equal, $k_x = k_y$, and the object is launched from point P in the $+y$ direction.</p>	 <p style="text-align: center;">$k_x = k_y$</p>
<p>b. The force constants are equal, $k_x = k_y$, and the object is launched from rest at point Q.</p>	 <p style="text-align: center;">$k_x = k_y$</p>

(continued on other side)


Pretest: Harmonic motion in two dimensions

- c. The force constants differ by a factor of 4, with $k_y = 4k_x$, and the object is launched from point R in the $+y$ direction.



$$k_y = 4k_x$$

- d. The force constants differ by a factor of 4, with $k_y = 4k_x$, and the object is launched from the origin in the direction shown.



$$k_y = 4k_x$$